

# Evidence Based Intervention

## Body Contouring

July 2022 v1.0

<b>Document Owner:</b>	Dr Rachel Joyce – Medical Director
<b>Document Author(s):</b>	Clinical Policies Group
<b>Version:</b>	v1.0
<b>Approved By:</b>	HWE ICB Commissioning Committee
<b>Date of Approval:</b>	1st July 2022
<b>Date of Review:</b>	1st July 2024



## Policy: Body Contouring

### Introduction

The aim of body contouring procedures is to improve the functional ability of people with severe loose or sagging skin. Body contouring is not funded for cosmetic or psychological reasons.

The areas where body contouring may be considered are (although this list is not exhaustive):

- Abdomen
- Thighs
- Arms - above the elbow only
- Breast

Abdominoplasty and apronectomy (body contouring of abdomen) are outside of the scope of this policy. Please refer instead to the national EBI programme criteria at <https://ebi.aomrc.org.uk/>  
This policy applies to body contouring at all other sites.

### Content

The body area that contouring is requested for must be made clear in the referral. Medical photography may be requested.

Body contouring should not be commissioned for cosmetic reasons and should be offered only in exceptional cases for those patients evidencing a strong clinical need.

The following criteria should be fulfilled:

- Age 18 or over  
AND
- Same site body contouring has not already been performed. (A maximum of 1 procedure for each area of the body will be funded.)  
AND
- Disfiguring scarring following trauma or previous surgery resulting in skin tethering to deep tissues and functional problems or severe pain.  
OR
- Excessive skin as a result of significant weight loss where:
  - Starting BMI above 40 and current BMI less than 25 with a stable weight maintained for at least 2 years.  
OR
  - Starting BMI above 50 and current BMI at least 20 points lower, or 50% of excess body weight lost, whichever is greater. Stable weight maintained for at least 2 years.

AND



- Significant functional problems
  - Experience of severe difficulty with activities of daily living as documented on the Barthel scale.  
OR
  - Significant recurrent infections beneath the skin fold, panniculitis or skin ulceration that has not responded to optimal medical therapy (typically for at least 12 months, requiring systemic antibiotics or antifungals)

All patients with excessive skin as a result of significant weight loss need a full assessment of their functional problems by an appropriate professional (e.g., Occupational Therapist) prior to an application, and the report must be included with the application. It is incumbent on the referring clinician only to refer for an assessment for patients who do have severe functional issues.

All funding for body contouring will be for a single stage procedure. Contour irregularities and moderate asymmetry are predictable following surgery. Any post-surgical cosmetic irregularities (including dog ears or unequal fat distribution) will not be funded by the ICB for revision surgery.

### Definitions

“Severe functional problems”: The following are examples of severe functional issues, but this list is not exhaustive:

- Documented evidence of recurrent infections beneath the skin folds
- Experiencing severe difficulties with daily living e.g., ambulatory restrictions, hygiene, ulcerations etc

“Stable weight”: Refers to weight with no greater fluctuation than +/- 5kg around the weight at the start of the two years. There must be evidence that weight independently assessed (e.g., through GP or weight management provider e.g., Weight Watchers/Slimming World) every 6 months during this time (i.e., 4 assessments taken over the 2 years).

“Excess body weight” is the weight loss required to reduce an individual to the maximum ‘normal’ BMI of 25kg/m<sup>2</sup>.

“Disfiguring scarring” – disfigurement is the state of having one’s appearance deeply and persistently harmed.

### Rationale

Procedures for aesthetic reasons are not routinely funded. Approval of body contouring is only for problems with severe functional problems and experiencing severe difficulties with the activities of daily living.

To make this procedure clinically effective, it is important that patients undergoing BCS procedures have achieved and maintained a stable weight so that the risks of recurrent obesity are reduced and risks relating to surgery are kept to a minimum.



**Patients who are not eligible for treatment under this policy may be considered on an individual basis where their GP or consultant believes exceptional circumstances exist that warrant deviation from the rule of this policy. Individual cases will be reviewed as per the ICB policy.**

## References

British Association of Plastic, Reconstructive and Aesthetic Surgeons, "UK Commissioning guide: Massive Weight Loss Body Contouring," 2017.

<http://www.bapras.org.uk/docs/defaultsource/commissioning-and-policy/2017--draft-for-consultation-body-contouring-surgerycommissioning.pdf?sfvrsn=0> (Accessed 19/10/18)

<http://www.bmj.com/company/wp-content/uploads/2016/05/NHS-weight-loss-surgery.pdf>

Al-Hadithy, H. Aditya, & K. Stewart, "Does the Degree of Ptosis Predict the Degree of Psychological Morbidity in Bariatric Patients Undergoing Reconstruction?" *Plastic and Reconstructive Surgery*, pp. 942-950, 2014.

Gilmartin, SJ, Long, A and Soldin, M, "Identity transformation and a changed lifestyle following dramatic weight loss and body-contouring surgery: An exploratory study," *Journal of Health Psychology*, vol. 20, no. 10, pp. 1318 - 1327, 2015.

van der Beek et al., "The impact of reconstructive procedures following bariatric surgery on patient wellbeing and quality of life," *Obesity Surgery*, vol. 2, no. 1, pp. 36-41, 2010.

de Zwaan et al., "Body image and quality of life in patients with and without body contouring surgery following bariatric surgery: a comparison of pre- and post- surgery groups," *Frontiers in Psychology*, vol. 5, pp. 1-10, 2014.




## Change History:

Version	Date	Reviewer(s)	Revision Description

## DOCUMENT CONTROL

*This is a controlled document. Whilst this document may be printed, the electronic version posted on the intranet is the controlled copy. Any printed copies of this document are not controlled. As a controlled document, this document should not be saved onto local or network drives but should always be accessed from the website.*

 *Do you really need to print this document?*

*Please consider the environment before you print this document and where copies should be printed double-sided. Please also consider setting the Page Range in the Print properties, when relevant to do so, to avoid printing the policy in its entirety.*

