



Hertfordshire and
West Essex Integrated
Care System



Hertfordshire and
West Essex
Integrated Care Board

Evidence Based Intervention

Shoulder Arthroscopy

Document Owner:	Dr Rachel Joyce – Medical Director
Document Author(s):	Clinical Policies Group
Version:	v1.1
Approved By:	Commissioning Committee
Date of Approval:	1 st July 2022
Date of Review:	July 2024



Policy: Shoulder Arthroscopy

Policy Summary

This policy covers the use of shoulder arthroscopy to treat a number of different conditions including labral tears, rotator cuff repair, adhesive capsulitis, and non-traumatic joint instability.

Arthroscopic shoulder decompression for subacromial shoulder pain is outside of the scope of this policy. Please refer instead to the national EBI policy at <https://ebi.aomrc.org.uk/> and the supplementary local policy on Arthroscopic shoulder decompression for subacromial shoulder pain.

Definition

An arthroscopy is a form of keyhole surgery that is used to look inside a joint and repair any damage that has occurred. An arthroscopy has two main uses:

- Treatment – an arthroscopy can be used to repair damage to the joint.
- Diagnosis – an arthroscopy can help diagnose problems with the joint, such as joint pain, stiffness, or limited range of joint movement.

The commissioners do not fund the use of shoulder arthroscopy for diagnostic purposes; radiological investigations should be used for this.

Threshold Eligibility Criteria

The patient will qualify for shoulder arthroscopy and treatment if clinically indicated, when they meet one of the following criteria:

- Full thickness rotator cuff tear as demonstrated by clinical symptoms and radiological imaging
OR
- Significant superior labrum anterior posterior (SLAP) tear as demonstrated by clinical symptoms and radiological imaging
OR
- Partial thickness rotator cuff tear as demonstrated by clinical symptoms and radiological imaging which has not responded to 3 months of conservative management
OR
- Minor (type I*) SLAP tear as demonstrated by clinical symptoms and radiological imaging which has not responded to 3 months of conservative management
OR
- *Adhesive capsulitis demonstrated by clinical symptoms which has not responded to 6 months of conservative management
OR
- *Adhesive capsulitis demonstrated by clinical symptoms and in the view of the treating consultant is having an extraordinarily severe impact on quality of life, and which has not responded to conservative management including corticosteroid injection where clinically appropriate.
OR
- Non-traumatic shoulder joint instability that has not responded to 6 months of conservative management
OR
- Traumatic shoulder joint instability alongside relevant conservative management as clinically appropriate.



Conservative Management

The conservative management to be attempted prior to referral includes the following:

- Activity modification
- Physiotherapy and exercise programme
- Oral analgesics, including NSAIDs (unless contraindicated)
- Steroid injections to the affected part of the joint where clinically appropriate

In the above criteria radiological imaging mentioned is to be organised by secondary care physicians as appropriate. Clinical symptoms are to be evaluated by both primary and secondary care physicians.

*Sydney classification (Synder SJ, Karzel RP, Del Pizzo W, et al. SLAP lesions of the shoulder. *Arthroscopy* 1990; 6; 274-279)

Frozen shoulders or adhesive capsulitis following a fracture WILL be funded as undertaking manipulation under anaesthetic increases the risk of a re-fracture.

Rationale

Rationale for shoulder arthroscopy includes adhesive capsulitis, rotator cuff damage and recurrent instability. In these cases, the evidence supports the use of shoulder arthroscopy for treatment purposes. However, the use of arthroscopy for diagnostic purposes is not supported and radiological investigations should be used for this.

In most circumstances, a clinical examination (history and examination) by a competent clinician will give a diagnosis and demonstrate if internal joint derangement is present. If there is diagnostic uncertainty despite competent examination or if there are “red flag” symptoms/signs/conditions, then an MRI scan might be indicated.

References

1. Coghlan JA, Buchbinder R, Green S, Johnston RV, Bell SN, Surgery for rotator cuff disease, *Cochrane Database of Systematic Reviews* 2008, Issue 1. Art. No.: CD005619. DOI: 10.1002/14651858.CD005619.pub2
2. NICE CKS revised April 2015 accessed online via <https://cks.nice.org.uk/shoulder-pain>
3. Woo Hyung Lee et al, Clinical Outcomes of Conservative Treatment and Arthroscopic Repair of Rotator Cuff Tears: A Retrospective Observational Study, *Ann Rehabil Med* 2016;40(2):252-262 pISSN: 2234-0645 eISSN: 2234-0653
<http://dx.doi.org/10.5535/arm.2016.40.2.252>
4. Baums et. al. Functional outcome and general health status in patients after arthroscopic release in adhesive capsulitis. *Knee Surg Sports Traumatol Arthrosc.* 2007 May; 15(5):638-44.
5. Snow M, Boutros I, Funk L. Posterior arthroscopic capsular release in frozen shoulder. *Arthroscopy.* 2009 Jan; 25(1):19-23.
6. Fernandes MR. Arthroscopic treatment of adhesive capsulitis of the shoulder with minimum follow up of six years. *Acta Ortop Bras.* 2015 Mar-Apr; 23(2): 85–89. doi: 1590/1413- 78522015230200613 PMID: PMC4813413
7. Wei Dong et. al. Treatments for Shoulder Impingement Syndrome. A PRISMA Systematic Review and Network Meta-Analysis, *Medicine*, Volume 94, Number 10, March 2015.
8. Bhatnagar A1, Bhonsle S2, Mehta S1. Correlation between MRI and Arthroscopy in Diagnosis of Shoulder Pathology. *J Clin Diagn Res.* 2016 Feb; 10(2):RC18-21
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4800609/>
9. Frozen Shoulder - Amar Rangan, Lorna Goodchild, Jo Gibson, Peter Brownson, Michael Thomas, Jonathan Rees, and Ro Kulkarni
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4935124/10>.<http://www.isrctn.com/ISRCTN48804508>
10. Beard JD et al. Arthroscopic subacromial decompression for subacromial shoulder pain (CSAW): a multicentre, pragmatic, parallel group, placebo-controlled, three-group, randomised surgical trial. *The Lancet* November 20, 2017 [http://dx.doi.org/10.1016/S0140-6736\(17\)32457-1](http://dx.doi.org/10.1016/S0140-6736(17)32457-1)



Patients who are not eligible for treatment under this policy may be considered on an individual basis where their GP or consultant believes exceptional circumstances exist that warrant deviation from the rule of this policy. Individual cases will be reviewed as per the ICB policy.

Change History:

Version	Date	Reviewer(s)	Revision Description
V1.1	November 2023	M Skerry	Removed inactive web link Removed reference to CCG

DOCUMENT CONTROL

This is a controlled document. Whilst this document may be printed, the electronic version posted on the intranet is the controlled copy. Any printed copies of this document are not controlled. As a controlled document, this document should not be saved onto local or network drives but should always be accessed from the website.

 *Do you really need to print this document?*

Please consider the environment before you print this document and where copies should be printed double-sided. Please also consider setting the Page Range in the Print properties, when relevant to do so, to avoid printing the policy in its entirety.

